

SSTR3 Rabbit mAb

货号: B30062

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	45kDa
实测分子量	70kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-87MG,LO2,Mouse brain,Rat brain,Rat spleen
细胞定位	Cell membrane,Multi-pass membrane protein
纯化	Affinity purification

抗原信息

抗原信息	Recombinant fusion protein corresponding to Human SSTR3 .
序列	RSERRVTRMVVAVVALFVLCWMPFYVLNIVNVVCPLPEEPAFFGLYFLVVALPYANSCANPILYGFLSYRFKQGFRRVLLRPS RRVRSQEPTVGPPEKTEE


靶点信息

7777	120	#IB	\Box
417	91ì.	育	T

This gene encodes a member of the somatostatin receptor protein family. Somatostatins are peptide hor mones that regulate diverse cellular functions such as neurotransmission, cell proliferation, and endocrin e signaling as well as inhibiting the release of many hormones and other secretory proteins. Somatostatin has two active forms of 14 and 28 amino acids. The biological effects of somatostatins are mediated by a family of G-protein coupled somatostatin receptors that are expressed in a tissue-specific manner. Som atostatin receptors form homodimers and heterodimers with other members of the superfamily as well as with other G-protein coupled receptors and receptor tyrosine kinases. This protein is functionally coupled to adenylyl cyclase. Alternate splicing results in multiple transcript variants.

基因ID	6753
基因名	SSTR3
Swiss	P32745
别名	SSTR3;SS-3-R;SS3-R;SS3R;SSR-28

产品验证



Western blot analysis of SSTR3 expressed in U-87MG,LO2,Mouse brain using SSTR3 Rabbit mAb at 1:10 00. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Bl ocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn