

# IAP2 Rabbit mAb

货号: B30115

## 产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC ICC IP
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 ICC: 1:50 - 1:200 IP: 1:20 - 1:50
理论分子量	68kDa
实测分子量	72kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HepG2,RAW264.7
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein corresponding to Human IAP2 .
序列	NWKRGDSPTEKHKKLYPSCRFVQSLNSVNNLEATSQPTFPSSVTNSTHSLLPGTENSGYFRGSYSNSPSNPVNSRANQD FS

## 靶点信息

研究背景	This gene encodes a member of the IAP family of proteins that inhibit apoptosis by binding to tumor necr osis factor receptor-associated factors TRAF1 and TRAF2, probably by interfering with activation of ICE-lik e proteases. The encoded protein inhibits apoptosis induced by serum deprivation but does not affect apoptosis resulting from exposure to menadione, a potent inducer of free radicals. It contains 3 baculovirus I AP repeats and a ring finger domain. Transcript variants encoding the same isoform have been identified.
------	---

基因ID	330
基因名	BIRC3
Swiss	Q13489
别名	BIRC3;AIP1;API2;CIAP2;HAIP1;HIAP1;MALT2;MIHC;RNF49;c-IAP2

## 产品验证



Western blot analysis of IAP2 expressed in HepG2,RAW264.7 using IAP2 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

### 实验步骤

访问官网浏览详情: www.ablybio.cn