

SOX9 Rabbit mAb

货号: B30118

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC IP FC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 IP: 1:20 - 1:50 FC: 1:20 - 1:50
理论分子量	70kDa
实测分子量	70kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HepG2,HeLa
细胞定位	Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human SOX9.
序列	HPDYKYQPRRRKSVKNGQAEAEATEQTHISPNAIFKALQADSPHSSSGMSEVHSPGEHSGSQGPPTPPTTPKTDVQP GKADLKREGRPLPEGGRQPPIDFRDVDIGELSSDVISNIETFDVNEFDQYLPP

靶点信息

研究背景	The protein encoded by this gene recognizes the sequence CCTTGAG along with other members of the H MG-box class DNA-binding proteins. It acts during chondrocyte differentiation and, with steroidogenic fact or 1, regulates transcription of the anti-Muellerian hormone (AMH) gene. Deficiencies lead to the skeletal malformation syndrome campomelic dysplasia, frequently with sex reversal. [provided by RefSeq, Jul 200 8]
------	--

基因ID	6662
基因名	SOX9
Swiss	P48436
别名	CMD1;CMPD1;SRA1;SRXX2;SRXY10;SOX9;SRY-box 9

产品验证



Western blot analysis of SOX9 expressed in HepG2,HeLa using SOX9 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn