

# CD116 Rabbit mAb

货号: B30161

## 产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	26kDa/31kDa/32kDa/38kDa/43kDa/46kDa/49kDa
实测分子量	50kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	bewo,293F(Negative control)
细胞定位	Cell membrane,Secreted,Single-pass type I membrane protein
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein corresponding to Human CD116.
序列	

## 靶点信息

研究背景	The protein encoded by this gene is the alpha subunit of the heterodimeric receptor for colony stimulatin g factor 2, a cytokine which controls the production, differentiation, and function of granulocytes and mac rophages. The encoded protein is a member of the cytokine family of receptors. This gene is found in the pseudoautosomal region (PAR) of the X and Y chromosomes. Multiple transcript variants encoding differe nt isoforms have been found for this gene, with some of the isoforms being membrane-bound and others being soluble.
基因 <b>ID</b>	1438

基因名	CSF2RA
Swiss	P15509
别名	CSF2RA;CD116;CDw116;CSF2R;CSF2RAX;CSF2RAY;CSF2RX;CSF2RY;GM-CSF-R-alpha;GMCSFR;GMR;SMDP 4

#### 产品验证



Western blot analysis of CD116 expressed in bewo,293F(Negative control) using CD116 Rabbit mAb at 1:1 000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blo cking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

#### 实验步骤

访问官网浏览详情: www.ablybio.cn