

STAT1 Rabbit mAb

货号: B30179

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB FC IP
推荐浓度	WB: 1:500 - 1:2000 FC: 1:20 - 1:50 IP: 1:20 - 1:50
理论分子量	83kDa/87kDa
实测分子量	84kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,Mouse spleen,Rat spleen
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human STAT1.
序列	NYKVMAAENIPENPLKYLYPNIDKDHAFGKYYSRPKEAPEPMELDGPKGTGYIKTELISVSEVHPSRLQTTDNLLPMSPEEFD EVSRIVGSVEFDSMMNTV

靶点信息

研究背景	The protein encoded by this gene is a member of the STAT protein family. In response to cytokines and gr owth factors, STAT family members are phosphorylated by the receptor associated kinases, and then for m homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. T his protein can be activated by various ligands including interferon-alpha, interferon-gamma, EGF, PDGF and IL6. This protein mediates the expression of a variety of genes, which is thought to be important for c ell viability in response to different cell stimuli and pathogens. Two alternatively spliced transcript variant s encoding distinct isoforms have been described.
基因ID	6772
基因名	STAT1
Swiss	P42224
别名	CANDF7;IMD31A;IMD31B;IMD31C;ISGF-3;STAT91;STAT1 alpha;STAT1

产品验证



Western blot analysis of STAT1 expressed in HeLa,Mouse spleen,Rat spleen using STAT1 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lan e. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn