

# Claudin 4 Rabbit mAb

货号: B30187

## 产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IP
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IP: 1:20 - 1:50
理论分子量	22kDa
实测分子量	18kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HT-29,A-431
细胞定位	Cell junction,Cell membrane,Multi-pass membrane protein,tight junction
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant fusion protein corresponding to Human Claudin 4 .
序列	VGGKCTNCLEDESAKAKTMIVAGVVFLLAGLMVIVPVSWTAHNIIQDFYNPLVASGQKREMGASLYVGWAASGLLLLGGG LLCCNCPPRTDKPYSAKYSAARSAAASNYV

## 靶点信息

研究背景	The protein encoded by this intronless gene belongs to the claudin family. Claudins are integral membran e proteins that are components of the epithelial cell tight junctions, which regulate movement of solutes and ions through the paracellular space. This protein is a high-affinity receptor for Clostridium perfringens enterotoxin (CPE) and may play a role in internal organ development and function during pre- and postna tal life. This gene is deleted in Williams-Beuren syndrome, a neurodevelopmental disorder affecting multi ple systems.
------	---

基因ID	1364
基因名	CLDN4
Swiss	O14493
别名	CLDN4;CPE-R;CPETR;CPETR1;WBSCR8;hCPE-R;claudin-4

## 产品验证



Western blot analysis of Claudin 4 expressed in HT-29,A-431 using Claudin 4 Rabbit mAb at 1:1000. Sec ondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking b uffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

### 实验步骤

访问官网浏览详情: www.ablybio.cn