

Synapsin II Rabbit mAb

货号: B30218

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200
理论分子量	63kDa
实测分子量	75kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T
细胞定位	glutamatergic synapse,plasma membrane,postsynaptic density,Schaffer collateral - CA1 synapse,SNARE complex,synapse,synaptic vesicle membrane
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human Synapsin II.
序列	HLIIGMQYAGLPSINSLESIYNFCDKPWVFAQLVAIYKTLGGEKFPLIEQTYYPNHKEMLTLPTFPVVVKIGHAHSGMGKVKV ENHYDFQDIASVVALTQT

靶点信息

研究背景	This gene is a member of the synapsin gene family. Synapsins encode neuronal phosphoproteins which a ssociate with the cytoplasmic surface of synaptic vesicles. Family members are characterized by common protein domains, and they are implicated in synaptogenesis and the modulation of neurotransmitter relea se, suggesting a potential role in several neuropsychiatric diseases. This member of the synapsin family e ncodes a neuron-specific phosphoprotein that selectively binds to small synaptic vesicles in the presynaptic nerve terminal. Polymorphisms in this gene are associated with abnormal presynaptic function and rel ated neuronal disorders, including autism, epilepsy, bipolar disorder and schizophrenia. Alternative splicing of this gene results in multiple transcript variants. The tissue inhibitor of metalloproteinase 4 gene is lo cated within an intron of this gene and is transcribed in the opposite direction. [provided by RefSeq, Feb 2 014]
基因ID	6854
基因名	SYN2
Swiss	Q92777
别名	SYNII

产品验证



Western blot analysis of Synapsin II expressed in 293T using Synapsin II Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn