

Alpha SNAP Rabbit mAb

货号: **B30274**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	33kDa
实测分子量	33kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HL-60,THP-1,HeLa,Jurkat,Mouse brain,Mouse lung,Mouse heart,Mouse pancreas,Rat liver
细胞定位	Membrane,Peripheral membrane protein
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human Alpha SNAP.
序列	MDNSGKEAEAMALLAEERKVKNSQSFFSGLFGGSSKIEEACEIYARAANMFKMAKNWSAAGNAFCQAAQLHLQLQSK HDAATCFVDAGNAFKKADPQEAINCLMRAIEIYTDMGFRFTIAAKHHISIAEIIYETELVDIEKAI AHYEQSADYYKGEESNSSAN KCLLKVAGYAALLEQYQKAI DIYEQVGTNAMDSPLLKYS AKDYFFKAALCHFCIDMLNAKLAVQKYEELFPAFSDSRECKLM KKLLEAHEEQNVDSYTESVKEYDSISRLDQWLTTMLLRIKKTIQGDEEDLR

靶点信息

研究背景	This gene encodes a member of the soluble NSF attachment protein (SNAP) family. SNAP proteins play a critical role in the docking and fusion of vesicles to target membranes as part of the 20S NSF-SNAP-SNARE complex. The encoded protein plays a role in the completion of membrane fusion by mediating the interaction of N-ethylmaleimide-sensitive factor (NSF) with the vesicle-associated and membrane-associated SNAP receptor (SNARE) complex, and stimulating the ATPase activity of NSF. Alternatively spliced transcript variants have been observed for this gene.
基因ID	8775
基因名	NAPA
Swiss	P54920
别名	NAPA;SNAPA

产品验证



Western blot analysis of Alpha SNAP expressed in Mouse brain,Rat liver,HL-60 using Alpha SNAP Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn