

FMO3 Rabbit mAb

货号: B30296

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	60kDa
实测分子量	56kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse liver
细胞定位	Endoplasmic reticulum membrane,Microsome membrane
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human FMO3.
序列	RFKHENYGLMPLNGVLRKEPVFNDELPASILCGIVSVKPNVKEFTETSAIFEDGTIFEGIDCVIFATGYSFAYPFLDESIIKSRN NEIILFKGVFPPLLEKSTIAVIGFVQSLGAAIPTVDLQSRWAAQVIKGTCTLPSMEDMMNDINEKMEKKRKWFGKSETIQTD YIVYMDELSSFIGAKPNIPWLFLTDPKLAMEVYFGPCSPYQFRLVGPGQWPGARNAILTQWDRSLKPMQTRVVGRLQKPC FFFHWLKLFAIPILLIAVFLVLT

靶点信息

研究背景	Flavin-containing monooxygenases (FMO) are an important class of drug-metabolizing enzymes that catal yze the NADPH-dependent oxygenation of various nitrogen-,sulfur-, and phosphorous-containing xenobiot ics such as therapeutic drugs, dietary compounds, pesticides, and other foreign compounds. The human FMO gene family is composed of 5 genes and multiple pseudogenes. FMO members have distinct develop mental- and tissue-specific expression patterns. The expression of this FMO3 gene, the major FMO expres sed in adult liver, can vary up to 20-fold between individuals. This inter-individual variation in FMO3 expression levels is likely to have significant effects on the rate at which xenobiotics are metabolised and, the refore, is of considerable interest to the pharmaceutical industry. This transmembrane protein localizes to the endoplasmic reticulum of many tissues. Alternative splicing of this gene results in multiple transcript variants encoding different isoforms. Mutations in this gene cause the disorder trimethylaminuria (TMAu) which is characterized by the accumulation and excretion of unmetabolized trimethylamine and a distinct ive body odor. In healthy individuals, trimethylamine is primarily converted to the non odorous trimethylamine N-oxide.
基因ID	2328
基因名	FMO3
Swiss	P31513
别名	FMO3;FMOII;TMAU;dJ127D3.1

产品验证



Western blot analysis of FMO3 expressed in Mouse liver using FMO3 Rabbit mAb at 1:1000. Secondary ant ibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% no n-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn