

CDKN2C Rabbit mAb

货号**: B30351**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC IP
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 IP: 1:20 - 1:50
理论分子量	15kDa
实测分子量	15kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,293T,NIH/3T3,Mouse lung,Mouse testis,Rat spleen,Rat kidney
细胞定位	cytoplasm,cytosol,nucleus
纯化	Affinity purification

抗原信息

抗原信息	Recombinant fusion protein corresponding to Human CDKN2C.	
序列	MAEPWGNELASAAARGDLEQLTSLLQNNVNVNAQNGFGRTALQVMKLGNPEIARRLLLRGANPDLKDRTGFAVIHDAAR AGFLDTLQTLLEFQADVNIEDNEGNLPLHLAAKEGHLRVVE	

靶点信息

研究背景	The protein encoded by this gene is a member of the INK4 family of cyclin-dependent kinase inhibitors. T his protein has been shown to interact with CDK4 or CDK6, and prevent the activation of the CDK kinases, thus function as a cell growth regulator that controls cell cycle G1 progression. Ectopic expression of this gene was shown to suppress the growth of human cells in a manner that appears to correlate with the pr esence of a wild-type RB1 function. Studies in the knockout mice suggested the roles of this gene in regul ating spermatogenesis, as well as in suppressing tumorigenesis. Two alternatively spliced transcript varia nts of this gene, which encode an identical protein, have been reported. [provided by RefSeq, Jul 2008]
基因ID	1031
基因名	CDKN2C
Swiss	P42773
别名	INK4C; p18; p18-INK4C

产品验证



Western blot analysis of CDKN2C expressed in Mouse lung,Rat spleen,HeLa using CDKN2C Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn