

USP13 Rabbit mAb

货号**: B30371**

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IF/ICC IP FC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200 IP: 1:20 - 1:50 FC: 1:20 - 1:50
理论分子量	90kDa/97kDa
实测分子量	97kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse heart,Mouse brain,Rat heart,Rat brain
细胞定位	cytosol,nucleoplasm,nucleus
纯化	Affinity purification

抗原信息

抗原信息	Recombinant fusion protein corresponding to Human USP13.	
序列	MHLKRHVREKVRGASGGALPKRRNSKIFLDLDTDDDLNSDDYEYEDEAKLVIFPDHYEIALPNIEELPALVTIACDAVLSSKS PYRKQDPDTWENELPVSKYANNLTQLDN	

靶点信息

研究背景	Deubiquitinase that mediates deubiquitination of target proteins such as BECN1, MITF, SKP2 and USP10 a nd is involved in various processes such as autophagy and endoplasmic reticulum-associated degradatio n (ERAD. Component of a regulatory loop that controls autophagy and p53/TP53 levels: mediates deubiq uitination of BECN1, a key regulator of autophagy, leading to stabilize the PIK3C3/VPS34-containing comp lexes. Also deubiquitinates USP10, an essential regulator of p53/TP53 stability. In turn, PIK3C3/VPS34-con taining complexes regulate USP13 stability, suggesting the existence of a regulatory system by which PIK 3C3/VPS34-containing complexes regulate p53/TP53 protein levels via USP10 and USP13. Recruited by nu clear UFD1 and mediates deubiquitination of SKP2, thereby regulating endoplasmic reticulum-associated degradation (ERAD. Also regulates ERAD through the deubiquitination of UBL4A a component of the BAG6 /BAT3 complex. Mediates stabilization of SIAH2 independently of deubiquitinase activity: binds ubiquitinat ed SIAH2 and acts by impairing SIAH2 autoubiquitination. Has a weak deubiquitinase activity in vitro and preferentially cleaves 'Lys-63'-linked polyubiquitin chains. In contrast to USP5, it is not able to mediate u nanchored polyubiquitin disassembly. Able to cleave ISG15 in vitro; however, additional experiments are required to confirm such data.
基因ID	8975
基因名	USP13
Swiss	Q92995
别名	USP13;ISOT3;IsoT-3

产品验证



Western blot analysis of USP13 expressed in Mouse heart, Mouse brain, Rat heart using USP13 Rabbit m Ab at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug p er lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s

实验步骤

访问官网浏览详情: www.ablybio.cn