

BMP15 Rabbit mAb

货号: B30393

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB FC
推荐浓度	WB: 1:500 - 1:2000 FC: 1:20 - 1:50
理论分子量	45kDa
实测分子量	50kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	OVCAR3,Mouse brain,Mouse kidney,Mouse liver,Rat brain
细胞定位	Secreted
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human BMP15.
序列	LTRFNLSCHVEPWVQKNPTNHFPSSEGDSKPSLMSNAWKEMDITQLVQQRFWNNKGHRILRLRFMCQQQKDSGGLEL WHGTSSLDIAFLLLYFNDTHKSIRKAKFLPRGMEEFMERESLLRRTRQADGISAEVTAASSSKHSGPENNQCSLHPFQISFR QLGWDHWIIAPPFYTPNYCKGTCLRVLRDGLNSPNHAIIQNLINQLVDQSVPBPSCPYKVYPISVLMIEANGSILYKEYEGMI AESCTCR

靶点信息

研究背景	This gene encodes a secreted ligand of the TGF-beta (transforming growth factor-beta) superfamily of proteins. Ligands of this family bind various TGF-beta receptors leading to recruitment and activation of SMA D family transcription factors that regulate gene expression. The encoded preprotein is proteolytically processed to generate subunits of a disulfide-linked homodimer, or alternatively, a heterodimer, with the related protein, growth differentiation factor 9 (GDF9). This protein plays a role in oocyte maturation and follicular development, through activation of granulosa cells. Defects in this gene are the cause of ovarian dysgenesis and are associated with premature ovarian failure.
基因ID	9210
基因名	BMP15
Swiss	O95972
别名	BMP15;GDF9B;ODG2;POF4

产品验证



Western blot analysis of BMP15 expressed in Mouse brain, Rat brain, OVCAR3 using BMP15 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn