

# FGFR1 Rabbit mAb

## 货号**: B30400**

## 产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200
理论分子量	6kDa/16kDa/33kDa/55-95kDa
实测分子量	140kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-251MG,Rat brain,Mouse brain
细胞定位	Cell membrane,Cytoplasm,Cytoplasmic vesicle,Nucleus,Single-pass type I membrane protein,cytosol
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein corresponding to Human FGFR1.
序列	IDYYKK

## 靶点信息

研究背景	The protein encoded by this gene is a member of the fibroblast growth factor receptor (FGFR) family, whe re amino acid sequence is highly conserved between members and throughout evolution. FGFR family m embers differ from one another in their ligand affinities and tissue distribution. A full-length representativ e protein consists of an extracellular region, composed of three immunoglobulin-like domains, a single hy drophobic membrane-spanning segment and a cytoplasmic tyrosine kinase domain. The extracellular por tion of the protein interacts with fibroblast growth factors, setting in motion a cascade of downstream sig nals, ultimately influencing mitogenesis and differentiation. This particular family member binds both acid ic and basic fibroblast growth factors and is involved in limb induction. Mutations in this gene have been associated with Pfeiffer syndrome, Jackson-Weiss syndrome, Antley-Bixler syndrome, osteoglophonic dysp lasia, and autosomal dominant Kallmann syndrome 2. Chromosomal aberrations involving this gene are a ssociated with stem cell myeloproliferative disorder and stem cell leukemia lymphoma syndrome. Alterna tively spliced variants which encode different protein isoforms have been described; however, not all vari ants have been fully characterized.
基因ID	2260
基因名	FGFR1
Swiss	P11362
别名	FGFR1;BFGFR;CD331;CEK;ECCL;FGFBR;FGFR-1;FLG;FLT-2;FLT2;HBGFR;HH2;HRTFDS;KAL2;N-SAM;OGD;bF GF-R-1

## 产品验证



Western blot analysis of FGFR1 expressed in U-251MG,Rat brain,Mouse brain using FGFR1 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per I ane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: www.ablybio.cn