

BAG2 Rabbit mAb

货号: B30440

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC IP FC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 IP: 1:20 - 1:50 FC: 1:20 - 1:50
理论分子量	24kDa
实测分子量	24kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	A-549,PC-3,Jurkat
细胞定位	axon,cytosol,dendrite,dendritic microtubule
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human BAG2.
序列	MAQAKINAKANEGRFRCSSSMADRSSRLLESLDQLELRVEALREAATAVEQEKEILLEMHSIQNSQDMRQISDGEREELNL TANRLMGRTLTVESVET

靶点信息

研究背景	BAG proteins compete with Hip for binding to the Hsc70/Hsp70 ATPase domain and promote substrate release. All the BAG proteins have an approximately 45-amino acid BAG domain near the C terminus but differ markedly in their N-terminal regions. The predicted BAG2 protein contains 211 amino acids. The BAG domains of BAG1, BAG2, and BAG3 interact specifically with the Hsc70 ATPase domain in vitro and in mammalian cells. All 3 proteins bind with high affinity to the ATPase domain of Hsc70 and inhibit its chaperone activity in a Hip-repressible manner. [provided by RefSeq, Jul 2008]
基因ID	9532
基因名	BAG2
Swiss	O95816
别名	BAG-2; dj41711.2

产品验证



实验步骤

访问官网浏览详情: www.ablybio.cn