

CCR2/CKR2 Rabbit mAb

货号: **B30481**

产品信息

| | |
|-------|---|
| 反应 | Human,Mouse |
| 宿主 | Rabbit |
| 克隆性 | Monoclonal |
| 预测反应 | |
| 应用 | IHC FC IP |
| 推荐浓度 | IHC: 1:50 - 1:200 FC: 1:20 - 1:50 IP: 1:20 - 1:50 |
| 理论分子量 | 41kDa |
| 实测分子量 | 42KDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | RAW264.7 |
| 细胞定位 | Cell membrane,Multi-pass membrane protein |
| 纯化 | Affinity purification |

抗原信息

| | |
|------|---|
| 抗原信息 | Recombinant fusion protein corresponding to Human CCR2/CKR2. |
| 序列 | KARTVTFGGVTSVITWLVAVFASVPGIIFTKCQKEDSVYVCGPYFPRGWNNFHTIMRNILGLVLPLLIMVICYSGILKTLLRCR NEKKRHRAVRVIFTIMI |

靶点信息

| | |
|------|--|
| 研究背景 | This gene encodes two isoforms of a receptor for monocyte chemoattractant protein-1, a chemokine which specifically mediates monocyte chemotaxis. Monocyte chemoattractant protein-1 is involved in monocyte infiltration in inflammatory diseases such as rheumatoid arthritis as well as in the inflammatory response against tumors. The receptors encoded by this gene mediate agonist-dependent calcium mobilization and inhibition of adenylyl cyclase. This gene is located in the chemokine receptor gene cluster region. Two alternatively spliced transcript variants are expressed by the gene. |
|------|--|

| | |
|-------|--|
| 基因ID | 729230 |
| 基因名 | CCR2 |
| Swiss | P41597 |
| 别名 | CCR2;CC-CKR-2;CCR-2;CCR2A;CCR2B;CD192;CKR2;CKR2A;CKR2B;CMKBR2;MCP-1-R;FLJ78302 |

产品验证

实验步骤

访问官网浏览详情: www.ablybio.cn