

5HT2C Receptor Rabbit mAb

货号: B30511

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	28kDa/51kDa
实测分子量	52kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	SH-SY5Y,A-431,U-87MG,Jurkat,B cells
细胞定位	Cell membrane,Multi-pass membrane protein
纯化	Affinity purification

抗原信息

抗原信息	Recombinant fusion protein corresponding to Human 5HT2C Receptor.
序列	


靶点信息

7"11"	120	dis.	\Box
41111	Hi.	Ħ	古

This gene encodes a seven-transmembrane G-protein-coupled receptor. The encoded protein responds to signaling through the neurotransmitter serotonin. The mRNA of this gene is subject to multiple RNA editin g events, where adenosine residues encoded by the genome are converted to inosines. RNA editing is predicted to alter the structure of the second intracellular loop, thereby generating alternate protein forms with decreased ability to interact with G proteins. Abnormalities in RNA editing of this gene have been detected in victims of suicide that suffer from depression. In addition, naturally-occuring variation in the promoter and 5' non-coding and coding regions of this gene may show statistically-significant association with mental illness and behavioral disorders. Alternative splicing results in multiple different transcript variants.

基因ID	3358
基因名	HTR2C
Swiss	P28335
别名	HTR2C;5-HT1C;5-HT2C;5-HTR2C;5HTR2C;HTR1C

产品验证



Western blot analysis of 5HT2C Receptor expressed in SH-SY5Y,A-431,U-87MG using 5HT2C Recept or Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/p roteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. E xposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn