

Phospho-AMPK alpha 1 (S496) Rabbit mAb

货号: B30522

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IF/ICC IP
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200 IP: 1:20 - 1:50
理论分子量	64kDa/65kDa
实测分子量	64kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse liver, Mouse brain, Rat liver
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human Phospho-AMPK alpha 1 (S496).
序列	AKSGT

靶点信息

The protein encoded by this gene belongs to the ser/thr protein kinase family. It is the catalytic subunit of the 5'-prime-AMP-activated protein kinase (AMPK). AMPK is a cellular energy sensor conserved in all euka ryotic cells. The kinase activity of AMPK is activated by the stimuli that increase the cellular AMP/ATP ratio . AMPK regulates the activities of a number of key metabolic enzymes through phosphorylation. It protect s cells from stresses that cause ATP depletion by switching off ATP-consuming biosynthetic pathways. Alt ernatively spliced transcript variants encoding distinct isoforms have been observed.

基因ID	5562
基因名	PRKAA1
Swiss	Q13131
别名	PRKAA1;AMPK;AMPKa1

产品验证



Western blot analysis of Phospho-AMPK alpha 1 (S496) expressed in Mouse liver, Mouse brain, Rat liver using Phospho-AMPK alpha 1 (S496) Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn