

Caspase-1 Rabbit mAb

货号: B30530

产品信息

反应	Human,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200
理论分子量	10kDa/29kDa/35kDa/42kDa/45kDa
实测分子量	45kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HepG2,Mouse testis
细胞定位	Cytoplasm
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human Caspase-1.
序列	VDVKKNLTASDMTTELEAFAHRPEHKTSDSTFLVFMSHGIREGICGKKHSEQVPDILQLNAIFNMLNTKNCPSLKDKPKVIII QACRGDSPGVVWFKDSVG

靶点信息

研究背景	This gene encodes a protein which is a member of the cysteine-aspartic acid protease (caspase) family. S equential activation of caspases plays a central role in the execution-phase of cell apoptosis. Caspases ex ist as inactive proenzymes which undergo proteolytic processing at conserved aspartic residues to produ ce 2 subunits, large and small, that dimerize to form the active enzyme. This gene was identified by its ab ility to proteolytically cleave and activate the inactive precursor of interleukin-1, a cytokine involved in the processes such as inflammation, septic shock, and wound healing. This gene has been shown to induce cell apoptosis and may function in various developmental stages. Studies of a similar gene in mouse sugg est a role in the pathogenesis of Huntington disease. Alternative splicing results in transcript variants encoding distinct isoforms.
基因 ID	834
基因名	CASP1
Swiss	P29466
别名	CASP1; ICE; IL1BC; P45; caspase-1

产品验证



Western blot analysis of Caspase-1 expressed in HepG2,Mouse testis using Caspase-1 Rabbit mAb at 1:1 000. Secondary antibody: HRP Goat Anti-Rabbit lgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Bl ocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn