

Cyclophilin A Rabbit mAb

货号: B30539

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200
理论分子量	11kDa/18kDa
实测分子量	18kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	K-562,Jurkat,HepG2,HeLa,NIH/3T3,Mouse lung,mouse spleen,rat spleen,mouse brain,rat brain
细胞定位	Cytoplasm,Secreted
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human Cyclophilin A.
序列	MANAGPNTNGSQFFICTAKTEWLDGKHVVFGKVKEGMNIVEAMERFGSRNGKTSKKITIADCGQLE

靶点信息

研究背景	This gene encodes a member of the peptidyl-prolyl cis-trans isomerase (PPIase) family. PPIases catalyze the cis-trans isomerization of proline imidic peptide bonds in oligopeptides and accelerate the folding of proteins. The encoded protein is a cyclosporin binding-protein and may play a role in cyclosporin A-mediated immunosuppression. The protein can also interact with several HIV proteins, including p55 gag, Vpr, and capsid protein, and has been shown to be necessary for the formation of infectious HIV virions. Multiple pseudogenes that map to different chromosomes have been reported.
基因ID	5478

基因名	PPIA
Swiss	P62937
别名	PPIA;CYPA;CYPH;HEL-S-69p

产品验证



Western blot analysis of Cyclophilin A expressed in Mouse lung,rat spleen,K-562 using Cyclophilin A R abbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn