

Legumain Rabbit mAb

货号: B30541

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	36,57kDa
实测分子量	36,57kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,HepG2,MCF7,Mouse brain,Mouse spleen,Mouse kidney
细胞定位	Lysosome
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human Legumain.
序列	ANPRESSYACYYDEKRSTYLGDWYSVNWMEDSDVEDLTKETLHKQYHLVKSHTNTSHVMQYGNKTISTMKVMQFQGM KRKASSPVPLPPVTHL

靶点信息

研究背景	This gene encodes a cysteine protease that has a strict specificity for hydrolysis of asparaginyl bonds. Thi s enzyme may be involved in the processing of bacterial peptides and endogenous proteins for MHC class II presentation in the lysosomal/endosomal systems. Enzyme activation is triggered by acidic pH and app ears to be autocatalytic. Protein expression occurs after monocytes differentiate into dendritic cells. A full y mature, active enzyme is produced following lipopolysaccharide expression in mature dendritic cells. O verexpression of this gene may be associated with the majority of solid tumor types. This gene has a pse udogene on chromosome 13. Several alternatively spliced transcript variants have been described, but the biological validity of only two has been determined. These two variants encode the same isoform. [provided by RefSeq, Jul 2008]
基因ID	5641
基因名	LGMN
Swiss	Q99538
别名	AEP; LGMN1; PRSC1

产品验证



Western blot analysis of Legumain expressed in Mouse brain,293T,HepG2 using Legumain Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn