

Cdc16/APC6 Rabbit mAb

货号: B30547

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200
理论分子量	72kDa
实测分子量	72kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	NIH/3T3
细胞定位	Cytoplasm,centrosome,cytoskeleton,microtubule organizing center,spindle
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human Cdc16/APC6.
序列	LLRFLFENKLKKYNKPSETVIPESVDGLQENLDVVVSLAERHYYNCDFKMCYKLTSVVMEKDPFHASCLP VHIGTLVELNKA NELFYLSHKLVDLYPSNPV

靶点信息

研究背景	The protein encoded by this gene functions as a protein ubiquitin ligase and is a component of the multiprotein APC complex. The APC complex is a cyclin degradation system that governs exit from mitosis by targeting cell cycle proteins for degradation by the 26S proteasome. Each component protein of the APC complex is highly conserved among eukaryotic organisms. This protein, and other APC complex proteins, contain a tetratricopeptide repeat (TPR) domain; a protein domain that is often involved in protein-protein interactions and the assembly of multiprotein complexes. Multiple alternatively spliced transcript variants, encoding distinct proteins, have been identified. [provided by RefSeq, Jan 2016]
------	--

基因ID	8881
基因名	CDC16
Swiss	Q13042
别名	ANAPC6; APC6; CDC16Hs; CUT9

产品验证



实验步骤

访问官网浏览详情: www.ablybio.cn