

# S100 Rabbit mAb

货号: B30549

### 产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IP FC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IP: 1:20 - 1:50 FC: 1:20 - 1:50
理论分子量	11kDa
实测分子量	11kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse lung,Mouse kidney,Mouse heart
细胞定位	Cytoplasm
纯化	Affinity purification


### 抗原信息

抗原信息	Recombinant fusion protein corresponding to Human S100.
序列	GSELETAMETLINVFHAHSGKEGDKYKLSKKELKELLQTELSGFLDAQKDVDAVDKVMKELDENGDGEVDFQEYVVLVA ALTVACNNFFWENS

靶点信息

研究背景	The protein encoded by this gene is a member of the S100 family of proteins containing 2 EF-hand calciu m-binding motifs. S100 proteins are localized in the cytoplasm and/or nucleus of a wide range of cells, an d involved in the regulation of a number of cellular processes such as cell cycle progression and differenti ation. S100 genes include at least 13 members which are located as a cluster on chromosome 1q21. This protein may function in stimulation of Ca2+-induced Ca2+ release, inhibition of microtubule assembly, a nd inhibition of protein kinase C-mediated phosphorylation. Reduced expression of this protein has been i mplicated in cardiomyopathies. [provided by RefSeq, Jul 2008]
基因ID	6271
基因名	S100A1
Swiss	P23297
别名	S100;S100-alpha;S100A;S100A1

### 产品验证



Western blot analysis of S100 expressed in Mouse lung, Mouse kidney, Mouse heart using S100 Rabbit m Ab at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug p er lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: www.ablybio.cn