

TIMP2 Rabbit mAb

货号: B30619

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB FC
推荐浓度	WB: 1:500 - 1:2000 FC: 1:20 - 1:50
理论分子量	24kDa
实测分子量	22kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	A-431,Mouse testis,Rat testis
细胞定位	Secreted
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human TIMP2.
序列	VSEKEVDSGNDIYGNPIKRIQYEIKQIKMFKGPEKDIEFIYTAPSSAVCGVSLDVGGKKEYLIAGKAEGDGKMHITLCDFIVP WDTLSTTQKKSLNHRYQM

靶点信息

This gene is a member of the TIMP gene family. The proteins encoded by this gene family are nature bitors of the matrix metalloproteinases, a group of peptidases involved in degradation of the extract matrix. In addition to an inhibitory role against metalloproteinases, the encoded protein has a unique among TIMP family members in its ability to directly suppress the proliferation of endothelial cells. A sult, the encoded protein may be critical to the maintenance of tissue homeostasis by suppressing oliferation of quiescent tissues in response to angiogenic factors, and by inhibiting protease activity sues undergoing remodelling of the extracellular matrix.	ellular ie role As a re the pr
--	---

基因ID	7077
基因名	TIMP2
Swiss	P16035
别名	TIMP2;CSC-21K;DDC8

产品验证



Western blot analysis of TIMP2 expressed in A-431, Mouse testis, Rat testis using TIMP2 Rabbit mAb at 1: 1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn