

UBE4B Rabbit mAb

货号: B30638

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	130kDa
实测分子量	130kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	PC-12,SKOV3,NIH/3T3,K-562,Mouse liver,Mouse brain,Rat liver,Rat brain
细胞定位	cytoplasm,nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human UBE4B.
序列	MEELSADEIRRRRLARLAGGQTSQPTTPLTSPQRENPPGPPIAASAPGPSQSLGLNVHNMTPATSPIGASGVAHRSQSSEG VSSLSSSPSNSLETQSQSL

靶点信息

研究背景	The modification of proteins with ubiquitin is an important cellular mechanism for targeting abnormal or s hort-lived proteins for degradation. Ubiquitination involves at least three classes of enzymes: ubiquitin-ac tivating enzymes, or E1s, ubiquitin-conjugating enzymes, or E2s, and ubiquitin-protein ligases, or E3s. Thi s gene encodes an additional conjugation factor, E4, which is involved in multiubiquitin chain assembly. T his gene is also the strongest candidate in the neuroblastoma tumor suppressor genes. Alternatively splic ed transcript variants encoding distinct isoforms have been found for this gene. [provided by RefSeq, Jul 2008]
------	--

基因ID	10277
基因名	UBE4B
Swiss	O95155
别名	E4; HDNB1; UBOX3; UFD2; UFD2A

产品验证



Western blot analysis of UBE4B expressed in Mouse liver,Rat liver,PC-12 using UBE4B Rabbit mAb at 1: 1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn