

TMPRSS2 Rabbit mAb

货号: B30658

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC FC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 FC: 1:20 - 1:50
理论分子量	54kDa
实测分子量	54kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HT-29,A-549,PC-3,Mouse kidney,Mouse stomach,Rat kidney
细胞定位	Cell membrane,Secreted,Single-pass type II membrane protein
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human TMPRSS2.
序列	MALNSGSPPAIGPYYENHGYQPENPYPAQPTVVPTVYEVHPAQYYPSPVPQYAPRVLTQASNPVVCTQPKSPSGTVCTSKT KKALCITLTLGTFLVGAAL

靶点信息

This gene encodes a protein that belongs to the serine protease family. The encoded protein contains a t ype II transmembrane domain, a receptor class A domain, a scavenger receptor cysteine-rich domain and a protease domain. Serine proteases are known to be involved in many physiological and pathological pr ocesses. This gene was demonstrated to be up-regulated by androgenic hormones in prostate cancer cell s and down-regulated in androgen-independent prostate cancer tissue. The protease domain of this prote in is thought to be cleaved and secreted into cell media after autocleavage. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Sep 2008]
7113
TMPRSS2
O15393
PP9284; PRSS10

产品验证



Western blot analysis of TMPRSS2 expressed in Mouse kidney,Rat kidney,HT-29 using TMPRSS2 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30u g per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn