

# FGFR3 Rabbit mAb

货号: B30700

## 产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC FC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 FC: 1:20 - 1:50
理论分子量	98kDa
实测分子量	88kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,Mouse brain,Mouse lung,Rat lung,Rat kidney
细胞定位	Cell membrane,Cytoplasmic vesicle,Endoplasmic reticulum,Secreted,Single-pass type I membrane protei n
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein corresponding to Human FGFR3.
序列	MGAPACALALCVAVAIVAGASSESLGTEQRVVGRAAEVPGPEPGQQEQLVFGSGDAVELSCPPPGGGPMGPTVWVKDG TGLVPSERVLVGPQRLQVLNAS

靶点信息

研究背景	This gene encodes a member of the fibroblast growth factor receptor (FGFR) family, with its amino acid s equence being highly conserved between members and among divergent species. FGFR family members differ from one another in their ligand affinities and tissue distribution. A full-length representative protein would consist of an extracellular region, composed of three immunoglobulin-like domains, a single hydrophobic membrane-spanning segment and a cytoplasmic tyrosine kinase domain. The extracellular portion of the protein interacts with fibroblast growth factors, setting in motion a cascade of downstream signals, ultimately influencing mitogenesis and differentiation. This particular family member binds acidic and basic fibroblast growth hormone and plays a role in bone development and maintenance. Mutations in this gene lead to craniosynostosis and multiple types of skeletal dysplasia. Three alternatively spliced trans cript variants that encode different protein isoforms have been described. [provided by RefSeq, Jul 2009]
基因ID	2261
基因名	FGFR3
Swiss	P22607
别名	ACH;CD333;CEK2;HSFGFR3EX;JTK4;FGFR3

## 产品验证



Western blot analysis of FGFR3 expressed in 293T, Mouse brain, Mouse lung using FGFR3 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lan e. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: www.ablybio.cn