

USP22 Rabbit mAb

货号: B30704

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IF/ICC IP
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200 IP: 1:20 - 1:50
理论分子量	58kDa/59kDa
实测分子量	60kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-87MG,293T
细胞定位	Nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein corresponding to Human USP22.
序列	TAEARKRKAKSCICHVCGVHLNRLHSCLYCVFFGCFTKKHIHEHAKAKRHNLAIDLMYGGIYCFQCQDYIYDKDMEIIAKEEQ RKAWKMQGVGEKFSTWEP

靶点信息

研究背景	Histone deubiquitinating component of the transcription regulatory histone acetylation (HAT complex SA GA. Catalyzes the deubiquitination of both histones H2A and H2B, thereby acting as a coactivator. Recruit ed to specific gene promoters by activators such as MYC, where it is required for transcription. Required f or nuclear receptor-mediated transactivation and cell cycle progression.
基因ID	23326

基因名	USP22
Swiss	Q9UPT9
别名	USP22;USP3L

产品验证



Western blot analysis of USP22 expressed in U-87MG,293T using USP22 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn