

CYP27A1 Rabbit mAb

货号: **B30722**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC FC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 FC: 1:20 - 1:50
理论分子量	60kDa
实测分子量	50kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,Mouse kidney,Rat liver
细胞定位	Mitochondrion membrane
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human CYP27A1.
序列	NQVSDMAQLFYFALEAICYILFEKRIGCLQRSIPEDTVTFVRSIGLMFQNSLYATFLPKWTRPVLPFWKRYLDGWNAIFSFG KKLIDEKLEDMEAQLQAAGPDGIQVSGYLHFLASGQLSPREAMGSLPELLMAGVDTTSNTLTWALYHLSKDPEIQEALHE EVLGVVPAGQVPQHKDFAHMPLLKAVLKETLRLYPVV

靶点信息

研究背景	This gene encodes a member of the cytochrome P450 superfamily of enzymes. The cytochrome P450 proteins are monooxygenases which catalyze many reactions involved in drug metabolism and synthesis of cholesterol, steroids and other lipids. This mitochondrial protein oxidizes cholesterol intermediates as part of the bile synthesis pathway. Since the conversion of cholesterol to bile acids is the major route for removing cholesterol from the body, this protein is important for overall cholesterol homeostasis. Mutations in this gene cause cerebrotendinous xanthomatosis, a rare autosomal recessive lipid storage disease.
基因ID	1593
基因名	CYP27A1
Swiss	Q02318
别名	CYP27A1;CP27;CTX;CYP27

产品验证



Western blot analysis of CYP27A1 expressed in 293T, Mouse kidney, Rat liver using CYP27A1 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120 s.

实验步骤

访问官网浏览详情: www.ablybio.cn