

ACVR1 Rabbit mAb

货号: B30729

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IP
推荐浓度	WB: 1:500 - 1:2000 IP: 1:20 - 1:50
理论分子量	57kDa
实测分子量	57kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,A-549,MCF7,Mouse brain,Rat testis,Rat brain
细胞定位	Membrane,Single-pass type I membrane protein
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human ACVR1.
序列	VLWEVARRMVSNQGVEDYKPPFYDVVPNDPSFEDMRKVVCVDQQRPNIPNRWFSPTLTSLAKLMKECWYQNPSARLTA LRIKKTLTKIDNSLDKLTDC

靶点信息

研究背景	Activins are dimeric growth and differentiation factors which belong to the transforming growth factor-beta (TGF-beta) superfamily of structurally related signaling proteins. Activins signal through a heteromeric complex of receptor serine kinases which include at least two type I (I and IB) and two type II (II and IIB) receptors. These receptors are all transmembrane proteins, composed of a ligand-binding extracellular domain with cysteine-rich region, a transmembrane domain, and a cytoplasmic domain with predicted serine/threonine specificity. Type I receptors are essential for signaling; and type II receptors are required for binding ligands and for expression of type I receptors. Type I and II receptors form a stable complex after ligand binding, resulting in phosphorylation of type I receptors by type II receptors. This gene encodes activin A type I receptor which signals a particular transcriptional response in concert with activin type II receptors. Mutations in this gene are associated with fibrodysplasia ossificans progressive.
基因ID	90
基因名	ACVR1
Swiss	Q04771
别名	ACVR1; ACTRI; ACVR1A; ACVRLK2; ALK2; FOP; SKR1; TSRI; activin receptor type-1

产品验证



实验步骤

访问官网浏览详情: www.ablybio.cn