

# Human IgG1 Rabbit mAb

# 货号**: B30748**

## 产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	52kDa
实测分子量	52kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	
细胞定位	blood microparticle,external side of plasma membrane,extracellular exosome,extracellular region,extrace llular space,immunoglobulin complex, circulating
纯化	Affinity purification

## 抗原信息

抗原信息	Recombinant fusion protein corresponding to Human Human IgG1.
序列	

### 靶点信息

研究背景	As a monomeric immunoglobulin that is predominately involved in the secondary antibody response and the only isotype that can pass through the human placenta, Immunoglobulin G (IgG) is synthesized and s ecreted by plasma B cells, and constitutes 75% of serum immunoglobulins in humans. IgG antibodies prot ect the body against the pathogens by agglutination and immobilization, complement activation, toxin ne utralization, as well as the antibody-dependent cell-mediated cytotoxicity (ADCC). IgG tetramer contains t wo heavy chains (50 kDa) and two light chains (25 kDa) linked by disulfide bonds, that is the two identic al halves form the Y-like shape. IgG is digested by pepsin proteolysis into Fab fragment (antigen-binding f ragment) and Fc fragment ("crystallizable" fragment). IgG1 is most abundant in serum among the four Ig G subclasses (IgG1, 2, 3 and 4) and binds to Fc receptors (FcγR ) on phagocytic cells with high affinity.
基因ID	3500
基因名	
Swiss	P01857
别名	

#### 产品验证



Western blot analysis of Human IgG1 expressed in HUVEC using Human IgG1 Rabbit mAb at 1:1000. Seco ndary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buff er: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

#### 实验步骤

访问官网浏览详情: www.ablybio.cn