

PKM2 Rabbit mAb

货号: B30751

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC FC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 FC: 1:20 - 1:50
理论分子量	56kDa/57kDa/58kDa
实测分子量	58kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-251MG,293T,HeLa,A-549
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human PKM2.
序列	SEAGTAFIQTQQLHAAMADTFLEHMCRLDIDSPPTARNTGIICTIGPASRSVETLKEMIKSGMNVARLNFSGHGTHEYHAETI KNVRTATESFASDPILYRPVAVALDTKGPEIRTGLIKSGTAEVELKKGATLKITLDNAYMEKCDENILWLDYKNICKVVEVG SKIYVDDGLISLQVKQKGADFLVTEVENGGSLGSKKGVNLPGAAV

靶点信息

研究背景	This gene encodes a protein involved in glycolysis. The encoded protein is a pyruvate kinase that catalyzes the transfer of a phosphoryl group from phosphoenolpyruvate to ADP, generating ATP and pyruvate. This protein has been shown to interact with thyroid hormone and may mediate cellular metabolic effects induced by thyroid hormones. This protein has been found to bind Opa protein, a bacterial outer membrane protein involved in gonococcal adherence to and invasion of human cells, suggesting a role of this protein in bacterial pathogenesis. Several alternatively spliced transcript variants encoding a few distinct isoforms have been reported.
基因ID	5315
基因名	PKM
Swiss	P14618
别名	CTHBP;HEL-S-30;OIP3;PK3;PKM2;TCB;THBP1;PKM

产品验证



实验步骤

访问官网浏览详情: www.ablybio.cn