

IL8 Rabbit mAb

货号: B30771

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	11kDa
实测分子量	11kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Recombinant Human IL-8 Protein
细胞定位	Secreted
纯化	Affinity purification

抗原信息

抗原信息	Recombinant fusion protein corresponding to Human IL8.
序列	MTSKLAVALLAAFLISAALCEGAVLPRSAKELRCQCIKTYSKPFHPKFIKELRVIESGPHCANTEIIVKLSDGRELCLDPKENWV QRVVEKFLKRAENS

靶点信息

研究背景	The protein encoded by this gene is a member of the CXC chemokine family. This chemokine is one of the major mediators of the inflammatory response. This chemokine is secreted by several cell types. It functions as a chemoattractant, and is also a potent angiogenic factor. This gene is believed to play a role in the pathogenesis of bronchiolitis, a common respiratory tract disease caused by viral infection. This gene a nd other ten members of the CXC chemokine gene family form a chemokine gene cluster in a region map ped to chromosome 4q.
基因ID	3576

基因名	CXCL8
Swiss	P10145
别名	CXCL8;GCP-1;GCP1;IL8;LECT;LUCT;LYNAP;MDNCF;MONAP;NAF;NAP-1;NAP1

产品验证



Western blot analysis of IL8 expressed in Recombinant Human IL-8 Protein using IL8 Rabbit mAb at 1:100 0. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocki ng buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn