

Tyrosinase Rabbit mAb

货号: B30775

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	42kDa/60kDa
实测分子量	70-80kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	Rat brain
细胞定位	Melanosome membrane, Single-pass type I membrane protein
纯化	Affinity purification

抗原信息

抗原信息	Recombinant fusion protein corresponding to Human Tyrosinase.
序列	FAHEAPAFPLPWHLFLLRWEQEIQKLTGDENFTIPYWDWRDAEKCDICTDEYMGQHPTNPNLLSPASFFSSWQIVCSRL EEYNSHQSLCNGTPEGPLRRN


靶点信息

研究背景	The enzyme encoded by this gene catalyzes the first 2 steps, and at least 1 subsequent step, in the conversion of tyrosine to melanin. The enzyme has both tyrosine hydroxylase and dopa oxidase catalytic activities, and requires copper for function. Mutations in this gene result in oculocutaneous albinism, and nonpathologic polymorphisms result in skin pigmentation variation. The human genome contains a pseudogene similar to the 3' half of this gene.
------	---

基因ID	7299
基因名	TYR
Swiss	P14679
别名	TYR;ATN;CMM8;OCA1;OCA1A;OCAIA;SHEP3

产品验证

Western blot analysis of Tyrosinase expressed in Rat brain using Tyrosinase Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.



实验步骤

访问官网浏览详情: www.ablybio.cn