

PAR4 Rabbit mAb

货号: B30826

产品信息

| 反应 | Human Maura Dat |
|-------|--|
| JXIM. | Human,Mouse,Rat |
| 宿主 | Rabbit |
| 克隆性 | Monoclonal |
| 预测反应 | |
| 应用 | WB |
| 推荐浓度 | WB: 1:500 - 1:2000 |
| 理论分子量 | 36kDa |
| 实测分子量 | 40kDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | BT-474,HT-29,SKOV3,SW480,HepG2 |
| 细胞定位 | Cytoplasm, Nucleus |
| 纯化 | Affinity purification |

抗原信息


| 抗原信息 | Recombinant fusion protein corresponding to Human PAR4. |
|------|---|
| 序列 | IGKLKEEIDLLNRDLDDIEDENEQLKQENKTLLKVVGQLTR |

靶点信息

| 研究背景 | The tumor suppressor WT1 represses and activates transcription. The protein encoded by this gene is a WT1-interacting protein that itself functions as a transcriptional repressor. It contains a putative leucine zi pper domain which interacts with the zinc finger DNA binding domain of WT1. This protein is specifically u pregulated during apoptosis of prostate cells. |
|------|---|
| 基因ID | 5074 |
| 基因名 | PAWR |

| Swiss | Q96IZ0 |
|-------|-----------------|
| 别名 | PAWR;PAR4;Par-4 |

产品验证



Western blot analysis of PAR4 expressed in BT-474,HT-29,SKOV3 using PAR4 Rabbit mAb at 1:1000. Se condary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blockin g buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn