

# **SUPT5H Rabbit mAb**

货号: B30855

### 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	155kDa
实测分子量	155kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa
细胞定位	Nucleus
纯化	Affinity purification


### 抗原信息

抗原信息	Recombinant fusion protein corresponding to Human SUPT5H.
序列	PHYGSQTPLHDGSRTPAQSGAWDPNNPNTPSRAEEEYEYAFDDEPTPSPQAYGGTPNPQTPGYPDPSSPQVNPQYNPQ TPGTPAMYNTDQFSPYAAPSPQG

靶点信息

研究背景	Component of the DRB sensitivity-inducing factor complex (DSIF complex, which regulates mRNA process ing and transcription elongation by RNA polymerase II. DSIF positively regulates mRNA capping by stimul ating the mRNA guanylyltransferase activity of RNGTT/CAP1A. DSIF also acts cooperatively with the negat ive elongation factor complex (NELF complex to enhance transcriptional pausing at sites proximal to the promoter. Transcriptional pausing may facilitate the assembly of an elongation competent RNA polymera se II complex. DSIF and NELF promote pausing by inhibition of the transcription elongation factor TFIIS/S-II . TFIIS/S-II binds to RNA polymerase II at transcription pause sites and stimulates the weak intrinsic nucle ase activity of the enzyme. Cleavage of blocked transcripts by RNA polymerase II promotes the resumption of transcription from the new 3' terminus and may allow repeated attempts at transcription through nat ural pause sites. DSIF can also positively regulate transcriptional elongation and is required for the efficie nt activation of transcriptional elongation by the HIV-1 nuclear transcriptional activator, Tat. DSIF acts to suppress transcriptional pausing in transcripts derived from the HIV-1 LTR and blocks premature release of HIV-1 transcripts at terminator sequences.
基因ID	6829
基因名	SUPT5H
Swiss	O00267
别名	SUPT5H; SPT5; SPT5H; Tat-CT1; transcription elongation factor SPT5

### 产品验证



Western blot analysis of SUPT5H expressed in HeLa using SUPT5H Rabbit mAb at 1:1000. Secondary antib ody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: www.ablybio.cn