

TXNRD2 Rabbit mAb

货号: B30873

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200
理论分子量	30kDa/46kDa/53kDa/56kDa
实测分子量	57kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,HepG2,Mouse spleen
细胞定位	Mitochondrion
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human TXNRD2.
序列	AALLGGLIQDAPNYGWEVAQPVPHPDWRKMAEAVQNHVKSLNWGHRVQLQDRKVKYFNIKASFVDEHTVCGVAKGGKEI LLSADHIIIATGGRPRYPTHIEGALEYGITSDDIFWLKESPGKTLVVGASYVALECAGFLTIGIGLDTTIMMRSIPLRGFDQQMS SMVIEHMASHGTRFLRGCAPSRVRRLLPDGQLQVTWEDSTTGKEDTGTFD

靶点信息

研究背景	The protein encoded by this gene belongs to the pyridine nucleotide-disulfide oxidoreductase family, and is a member of the thioredoxin (Trx) system. Three thioredoxin reductase (TrxR) isozymes are found in mammals. TrxRs are selenocysteine-containing flavoenzymes, which reduce thioredoxins, as well as other substrates, and play a key role in redox homoeostasis. This gene encodes a mitochondrial form important for scavenging reactive oxygen species in mitochondria. It functions as a homodimer containing FAD, and selenocysteine (Sec) at the active site. Sec is encoded by UGA codon that normally signals translation termination. The 3' UTRs of selenoprotein mRNAs contain a conserved stem-loop structure, the Sec insertion sequence (SECIS) element, which is necessary for the recognition of UGA as a Sec codon rather than as a stop signal. Alternatively spliced transcript variants encoding different isoforms, including a few localized in the cytosol and some lacking the C-terminal Sec residue, have been found for this gene.
基因ID	10587
基因名	TXNRD2
Swiss	Q9NNW7
别名	TXNRD2;SELZ;TR;TR-BETA;TR3;TRXR2

产品验证



实验步骤

访问官网浏览详情: www.ablybio.cn