

Cdk4 Rabbit mAb

货号: B30874

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IF/ICC IP
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200 IP: 1:20 - 1:50
理论分子量	20kDa/33kDa
实测分子量	34kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,HeLa
细胞定位	Cytoplasm,Membrane,Nucleus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human Cdk4.
序列	VGCIFAEMFRRKPLFCGNSEADQLGKIFDLIGLPPEDDWPRDVSLPRGAFPPRGPRPVQSVVPEMEESGAQLLLEMLTFNP HKRISAFRALQHSYLHKDEGNPE

靶点信息

研究背景	The protein encoded by this gene is a member of the Ser/Thr protein kinase family. This protein is highly similar to the gene products of S. cerevisiae cdc28 and S. pombe cdc2. It is a catalytic subunit of the prot ein kinase complex that is important for cell cycle G1 phase progression. The activity of this kinase is rest ricted to the G1-S phase, which is controlled by the regulatory subunits D-type cyclins and CDK inhibitor p 16(INK4a). This kinase was shown to be responsible for the phosphorylation of retinoblastoma gene produ ct (Rb). Mutations in this gene as well as in its related proteins including D-type cyclins, p16(INK4a) and R b were all found to be associated with tumorigenesis of a variety of cancers. Multiple polyadenylation site s of this gene have been reported.
基因 ID	1019
基因名	CDK4
Swiss	P11802
别名	CDK4;CMM3;PSK-J3

产品验证



Western blot analysis of Cdk4 expressed in 293T,HeLa using Cdk4 Rabbit mAb at 1:1000. Secondary antib ody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn