

# ATF7 Rabbit mAb

货号: B30881

### 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	13kDa/33kDa/49kDa/50kDa/51kDa/52kDa
实测分子量	53kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	BT-474,SW620,HepG2
细胞定位	Cytoplasm, Nucleus, nucleoplasm
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein corresponding to Human ATF7.
序列	MGDDRPFVCNAPGCGQRFTNEDHLAVHKHKHEMTLKFGPARTDSVIIADQTPTPTRFLKNCEEVGLFNELASSFEHEFKK AADEDEKKARSRTVAKKLVVFRPRLFLLCFGIIFLIG

靶点信息

研究背景	Stress-responsive chromatin regulator that plays a role in various biological processes including innate im munological memory, adipocyte differentiation or telomerase regulation. In absence of stress, contributes to the formation of heterochromatin and heterochromatin-like structure by recruiting histone H3K9 tri- an d di-methyltransferases thus silencing the transcription of target genes such as STAT1 in adipocytes, or g enes involved in innate immunity in macrophages and adipocytes (By similarity. Stress induces ATF7 pho sphorylation that disrupts interactions with histone methyltransferase and enhances the association with coactivators containing histone acetyltransferase and/or histone demethylase, leading to disruption of the heterochromatin-like structure and subsequently transcriptional activation (By similarity. In response to TNF-alpha, which is induced by various stresses, phosphorylated ATF7 and telomerase are released from t elomeres leading to telomere shortening. Plays also a role in maintaining epithelial regenerative capacity and protecting against cell death during intestinal epithelial damage and repair (By similarity.
基因 <b>ID</b>	11016
基因名	ATF7
Swiss	P17544
别名	ATF7;ATFA

#### 产品验证



Western blot analysis of ATF7 expressed in BT-474,SW620,HepG2 using ATF7 Rabbit mAb at 1:1000. Se condary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: www.ablybio.cn