

Gli3 Rabbit mAb

货号: B30886

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200
理论分子量	190kDa
实测分子量	190kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa
细胞定位	axoneme, ciliary base, ciliary tip, cilium, cytoplasm, cytosol, GLI-SUFU complex, nuclear speck, nucleolus, nucleoplasm, nucleus
纯化	Affinity purification

抗原信息

抗原信息	Recombinant fusion protein corresponding to Human Gli3.
序列	PNESAGSMVNGMQNQDPVGQGYLAHQLLGDSMQHPGAGRPGQQMLGQISATSHINIYQGPESCLPGAHGMGSQPSSL AVVRGYQPCASFGGSRRQAMPRDSLALQSGQLSDTSQTCRVNGIKMEMKGQPHPLCSNLQNYSGQFYDQTVGFSQQD TKAGSFSISDASCLLQGTSAKNSELLSPGANQVTSTVDSLDSHDLEGVQIDFDAIIDDGDHSSLMSGALSPSIIQNLSHSSS RLTTPRASLPFPALSMSTTNMAIGDMSSLLTSLAEESKFLAVMQ

靶点信息

研究背景	This gene encodes a protein which belongs to the C2H2-type zinc finger proteins subclass of the Gli famil y. They are characterized as DNA-binding transcription factors and are mediators of Sonic hedgehog (Shh) signaling. The protein encoded by this gene localizes in the cytoplasm and activates patched Drosophila homolog (PTCH) gene expression. It is also thought to play a role during embryogenesis. Mutations in this gene have been associated with several diseases, including Greig cephalopolysyndactyly syndrome, Pallis ter-Hall syndrome, preaxial polydactyly type IV, and postaxial polydactyly types A1 and B. [provided by R efSeq, Jul 2008]
基因ID	2737
基因名	GLI3
Swiss	P10071
别名	ACLS; GCPS; GLI3-190; GLI3FL; PAP-A; PAPA; PAPA1; PAPB; PHS; PPDIV

产品验证



Western blot analysis of Gli3 expressed in HeLa using Gli3 Rabbit mAb at 1:1000. Secondary antibody: HR P Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn