

# **PIAS2 Rabbit mAb**

货号: B30899

# 产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200
理论分子量	45kDa/63kDa/68kDa
实测分子量	55kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse testis
细胞定位	Nucleus,Nucleus speckle,PML body
纯化	Affinity purification


# 抗原信息

抗原信息	Recombinant fusion protein corresponding to Human PIAS2.
序列	EDGSWCPMRPKKEAMKVSSQPCTKIESSSVLSKPCSVTVASEASKKKVDVIDLTIESSSDEEEDPPAKRKCIFMSETQSSP TKGVLMYQPSSVRVPSVTSVDPAAIPPSLTDYSVPFHHTPISSMSSDLPGLDFLSLIPVDPQYCPPMFLDSLTSPLTASSTSV TTTSSHESSTHVSSSSSRSETGVITSSGSNIPDIISLD

靶点信息

研究背景	This gene encodes a member of the protein inhibitor of activated STAT (PIAS) family. PIAS proteins functio n as SUMO E3 ligases and play important roles in many cellular processes by mediating the sumoylation of target proteins. Alternatively spliced transcript variants encoding multiple isoforms have been observe d for this gene. Isoforms of the encoded protein enhance the sumoylation of specific target proteins including the p53 tumor suppressor protein, c-Jun, and the androgen receptor. A pseudogene of this gene is located on the short arm of chromosome 4. The symbol MIZ1 has also been associated with ZBTB17 which is a different gene located on chromosome 1.
基因ID	9063
基因名	PIAS2
Swiss	O75928
别名	PIAS2;ARIP3;DIP;MIZ;MIZ1;PIASX;PIASX-ALPHA;PIASX-BETA;SIZ2;ZMIZ4

### 产品验证



Western blot analysis of PIAS2 expressed in Mouse testis using PIAS2 Rabbit mAb at 1:1000. Secondary a ntibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

### 实验步骤

访问官网浏览详情: www.ablybio.cn