

Semaphorin 4D / CD100 Rabbit mAb

货号: B30916

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	82kDa/96kDa
实测分子量	120,140kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Daudi,PC-3
细胞定位	Cell membrane,Single-pass type I membrane protein
纯化	Affinity purification

抗原信息

抗原信息	Recombinant fusion protein corresponding to Human Semaphorin 4D / CD100.
序列	


靶点信息

7"11"	120	dis.	\Box
41111	Hi.	Ħ	古

This protein is a member of the semaphorin family of proteins and an important mediator of the moveme nt and differentiation of multiple cell types, including those of the immune, vascular, and nervous system s. VEGF and SEMA4D had a positive correlation with the malignant degree of ovarian cancer, and SEMA4D can serve as an independent prognostic factor. SEMA4D was the first semaphorin described to have im mune functions and serves important roles in T cell priming, antibody production, and cell-to-cell adhesio n. Proteolytic cleavage of SEMA4D from the cell surface gives rise to a soluble fragment of SEMA4D (sSEM A4D). Similar to the transmembranal form, sSEMA4D is thought to have immunoregulatory properties.

基因ID	10507
基因名	SEMA4D
Swiss	Q92854
别名	SEMA4D;C9orf164;CD100;M-sema-G;SEMAJ;coll-4

产品验证



Western blot analysis of Semaphorin 4D / CD100 expressed in Daudi,PC-3 using Semaphorin 4D / CD100 Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit lgG (H+L) at 1:5000. Lysat es/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn