

# **CDK1** Rabbit mAb

货号: B30936

## 产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC IP
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 IP: 1:20 - 1:50
理论分子量	27kDa/34kDa
实测分子量	34kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HT-29,HeLa,NIH/3T3,C6
细胞定位	Cytoplasm,Mitochondrion,Nucleus,centrosome,cytoskeleton,microtubule organizing center,spindle
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein corresponding to Human CDK1.
序列	GTYGV

靶点信息

研究背景	The protein encoded by this gene is a member of the Ser/Thr protein kinase family. This protein is a catal ytic subunit of the highly conserved protein kinase complex known as M-phase promoting factor (MPF), w hich is essential for G1/S and G2/M phase transitions of eukaryotic cell cycle. Mitotic cyclins stably associ ate with this protein and function as regulatory subunits. The kinase activity of this protein is controlled by cyclin accumulation and destruction through the cell cycle. The phosphorylation and dephosphorylation of this protein also play important regulatory roles in cell cycle control. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.
基因ID	983
基因名	CDK1
Swiss	P06493
别名	CDK1;CDC2;CDC28A;P34CDC2

#### 产品验证



Western blot analysis of CDK1 expressed in HT-29,HeLa,NIH/3T3 using CDK1 Rabbit mAb at 1:1000. Se condary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

#### 实验步骤

访问官网浏览详情: www.ablybio.cn