

DLDH Rabbit mAb

货号: B30955

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	56kDa
实测分子量	56kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	SKOV3,U-251MG,Mouse liver,Mouse brain,Rat brain
细胞定位	Mitochondrion matrix
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human DLDH.
序列	WVGKSEEQLKEEGIEYKVGKFPFAANSRAKTNADTDGMVKILGQKSTDRVLGAHILGPGAGEMVNEAALALEYGASCEDI ARVCHAHPTLSEAFREANLA

靶点信息

研究背景	This gene encodes a member of the class-I pyridine nucleotide-disulfide oxidoreductase family. The encod ed protein has been identified as a moonlighting protein based on its ability to perform mechanistically di stinct functions. In homodimeric form, the encoded protein functions as a dehydrogenase and is found in several multi-enzyme complexes that regulate energy metabolism. However, as a monomer, this protein can function as a protease. Mutations in this gene have been identified in patients with E3-deficient mapl e syrup urine disease and lipoamide dehydrogenase deficiency. Alternative splicing results in multiple tra nscript variants. [provided by RefSeq, Jan 2014]
基因ID	1738
基因名	DLD
Swiss	P09622
别名	DLDD; DLDH; E3; GCSL; LAD; PHE3

产品验证



Western blot analysis of DLDH expressed in Mouse liver,Rat brain,SKOV3 using DLDH Rabbit mAb at 1:1 000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. B locking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn