

# Phospho-Nucleophosmin (T199) Rabbit mAb

货号: B30981

#### 产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC FC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 FC: 1:20 - 1:50
理论分子量	28kDa/29kDa/32kDa
实测分子量	33kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,PC-12
细胞定位	Cytoplasm, Nucleus, centrosome, cytoskeleton, microtubule organizing center, nucleolus, nucleoplasm
纯化	Affinity purification

# 抗原信息


抗原信息	Recombinant fusion protein corresponding to Human Phospho-Nucleophosmin (T199).
序列	RDTPA

### 靶点信息

研究背景	This gene encodes a phosphoprotein which moves between the nucleus and the cytoplasm. The gene pro duct is thought to be involved in several processes including regulation of the ARF/p53 pathway. A numb er of genes are fusion partners have been characterized, in particular the anaplastic lymphoma kinase ge ne on chromosome 2. Mutations in this gene are associated with acute myeloid leukemia. More than a do zen pseudogenes of this gene have been identified. Alternative splicing results in multiple transcript varia nts.
------	---

基因ID	4869
基因名	NPM1
Swiss	P06748
别名	B23;NPM;NPM1

# 产品验证



Western blot analysis of Phospho-Nucleophosmin (T199) expressed in HeLa,PC-12 using Phospho-Nucleophosmin (T199) Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Det ection: ECL Enhanced Kit. Exposure time: 120s.

### 实验步骤

访问官网浏览详情: www.ablybio.cn