

Phospho-LRRK2 (S935) Rabbit mAb

货号: B31038

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200
理论分子量	286kDa
实测分子量	286kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Rat lung
细胞定位	Cell projection,Cytoplasm,Cytoplasmic side,Cytoplasmic vesicle,Endoplasmic reticulum,Endosome,Golgi apparatus,Lysosome,Membrane,Mitochondrion,Mitochondrion inner membrane,Mitochondrion matrix,Mitochondrion outer membrane,Perikaryon,Peripheral membrane protein,axon,dendrite,secretory vesicle,synaptic vesicle membrane
纯化	Affinity purification

抗原信息

抗原信息	Recombinant fusion protein corresponding to Human Phospho-LRRK2 (S935) .
序列	EKESSPKLVELLNNSGSREQDVRKALTISIGKGDSQISSLRLALDVANNNSICLGGFCIGKVEPSWLGPLFPDKTSNLRQTNIASTLARMVIRYQMKSVEEGTAGSDGNFSEDVLSKFDEWTFIPDSSMDSVFAQSDDLDSEGSEGSFLVKKKSNS


靶点信息

研究背景	This gene is a member of the leucine-rich repeat kinase family and encodes a protein with an ankyrin repeat region, a leucine-rich repeat (LRR) domain, a kinase domain, a DFG-like motif, a RAS domain, a GTPase domain, a MLK-like domain, and a WD40 domain. The protein is present largely in the cytoplasm but also associates with the mitochondrial outer membrane. Mutations in this gene have been associated with Parkinson disease-8.
------	--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

基因ID	120892
基因名	LRRK2
Swiss	Q5S007
别名	LRRK2;AURA17;DARDARIN;PARK8;RIPK7;ROCO2

产品验证

Western blot analysis of Phospho-LRRK2 (S935) expressed in Rat lung using Phospho-LRRK2 (S935) Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.



实验步骤

访问官网浏览详情: www.ablybio.cn