

Smad1 Rabbit mAb

货号: B31070

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200
理论分子量	15kDa/52kDa
实测分子量	60kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	BMP2 treated HepG2 cells
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human Smad1.
序列	

靶点信息

研究背景	The protein encoded by this gene belongs to the SMAD, a family of proteins similar to the gene products of the Drosophila gene 'mothers against decapentaplegic' (Mad) and the C. elegans gene Sma. SMAD prot eins are signal transducers and transcriptional modulators that mediate multiple signaling pathways. This protein mediates the signals of the bone morphogenetic proteins (BMPs), which are involved in a range of biological activities including cell growth, apoptosis, morphogenesis, development and immune response s. In response to BMP ligands, this protein can be phosphorylated and activated by the BMP receptor kina se. The phosphorylated form of this protein forms a complex with SMAD4, which is important for its functi on in the transcription regulation. This protein is a target for SMAD-specific E3 ubiquitin ligases, such as S MURF1 and SMURF2, and undergoes ubiquitination and proteasome-mediated degradation. Alternatively spliced transcript variants encoding the same protein have been observed.
基因 ID	4086
基因名	SMAD1
Swiss	Q15797
别名	SMAD1; BSP-1; BSP1; JV4-1; JV41; MADH1; MADR1; SMAD family member 1

产品验证



Western blot analysis of Smad1 expressed in BMP2 treated HepG2 cells using Smad1 Rabbit mAb at 1:100 0. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocki ng buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn