

# NgR3 Rabbit mAb

货号: B31090

## 产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	49kDa
实测分子量	49kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	SH-SY5Y,Mouse brain,Rat brain,T47D
细胞定位	Cell membrane,GPI-anchor,Lipid-anchor
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein corresponding to Human NgR3.
序列	PHGPRPGHRKPGKNCTNPRNRNQISKAGAGKQAPELPDYAPDYQHKFSFDIMPTARPKRKGKCARRTPIRAPSGVQQAS SASSLGASLLAWTLGLAVTLR

靶点信息

研究背景	Cell surface receptor. Plays a functionally redundant role in postnatal brain development and in regulatin g axon regeneration in the adult central nervous system. Contributes to normal axon migration across the brain midline and normal formation of the corpus callosum. Protects motoneurons against apoptosis; protection against apoptosis is probably mediated by MAG. Plays a role in inhibiting neurite outgrowth and axon regeneration via its binding to neuronal chondroitin sulfate proteoglycans. Binds heparin (By similar ity. Like other family members, plays a role in restricting the number dendritic spines and the number of synapses that are formed during brain development. Signaling mediates activation of Rho and downstrea m reorganization of the actin cytoskeleton.
基因 <b>ID</b>	146760
基因名	RTN4RL1
Swiss	Q86UN2
别名	RTN4RL1;NGRH2;NgR3

## 产品验证



Western blot analysis of NgR3 expressed in SH-SY5Y,Mouse brain,Rat brain using NgR3 Rabbit mAb at 1 :1000. Secondary antibody: HRP Goat Anti-Rabbit lgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

### 实验步骤

访问官网浏览详情: www.ablybio.cn