

ZBTB7A Rabbit mAb

货号: B31132

产品信息

| 反应 | Human |
|-------|--|
| 宿主 | Rabbit |
| 克隆性 | Monoclonal |
| 预测反应 | |
| 应用 | WB IHC IF/ICC IP |
| 推荐浓度 | WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 IP: 1:20 - 1:50 |
| 理论分子量 | 70kDa |
| 实测分子量 | 70kDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | HT-29,MCF7 |
| 细胞定位 | cytoplasm,nucleus |
| 纯化 | Affinity purification |


抗原信息

| 抗原信息 | Recombinant fusion protein corresponding to Human ZBTB7A. |
|------|---|
| 序列 | NYDLKNHMRVHTGLRPYQCDSCCKTFVRSDHLHRHLKKDGCNGVPSRRGRKPRVRGGAPDPSPGATATPGAPAQPSSP DARRNGQEKHFKDEDEDEDVASP |

靶点信息

| 研究背景 | Transcription factor that represses the transcription of a wide range of genes involved in cell proliferation and differentiation. Directly and specifically binds to the consensus sequence 5'-[GA][CA]GACCCCCCCC-3' and represses transcription both by regulating the organization of chromatin and through the direct rec ruitment of transcription factors to gene regulatory regions. Negatively regulates SMAD4 transcriptional a ctivity in the TGF-beta signaling pathway through these two mechanisms. That is, recruits the chromatin r egulator HDAC1 to the SMAD4-DNA complex and in parallel prevents the recruitment of the transcriptional activators CREBBP and EP300. Collaborates with transcription factors like RELA to modify the accessibility of gene transcription regulatory regions to secondary transcription factors (By similarity. Also directly in teracts with transcription factors like SP1 to prevent their binding to DNA. Functions as an androgen receptor/AR transcriptional corepressor by recruiting NCOR1 and NCOR2 to the androgen response elements/ARE on target genes. Thereby, negatively regulates androgen receptor signaling and androgen-induced cell proliferation. |
|--------------|--|
| 基因 ID | 51341 |
| 基因名 | ZBTB7A |
| Swiss | O95365 |
| 别名 | FBI-1; FBI1; LRF; TIP21; ZBTB7; ZNF857A; pokemon |

产品验证



Western blot analysis of ZBTB7A expressed in HT-29,MCF7 using ZBTB7A Rabbit mAb at 1:1000. Second ary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffe r: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn