

NOTCH4 Rabbit mAb

货号: B31147

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB FC
推荐浓度	WB: 1:500 - 1:2000 FC: 1:20 - 1:50
理论分子量	39kDa/61kDa/209kDa
实测分子量	230kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse kidney
细胞定位	Cell membrane,Nucleus,Single-pass type I membrane protein
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human NOTCH4.
序列	PPEARHKATPGREAGPFPRARTVSVSVPPHGGGALPRCRTLSAGAGPRGGGACLQARTWSVDLAARGGGAYSHCRSLS GVGAGGGPTPRGRRFSAGMRGPRPNPAIMRGRYGVAAGRGGRVSTDDWPCDWVALGACGSASNIPIPPCLTPSPERG SPQLDCGPPALQEMPINQGEGKK

靶点信息

研究背景	This gene encodes a member of the NOTCH family of proteins. Members of this Type I transmembrane protein family share structural characteristics including an extracellular domain consisting of multiple epidermal growth factor-like (EGF) repeats, and an intracellular domain consisting of multiple different domain types. Notch signaling is an evolutionarily conserved intercellular signaling pathway that regulates interactions between physically adjacent cells through binding of Notch family receptors to their cognate ligands . The encoded preproprotein is proteolytically processed in the trans-Golgi network to generate two polypeptide chains that heterodimerize to form the mature cell-surface receptor. This receptor may play a role in vascular, renal and hepatic development. Mutations in this gene may be associated with schizophrenia . Alternative splicing results in multiple transcript variants, at least one of which encodes an isoform that is proteolytically processed.
基因ID	4855
基因名	NOTCH4
Swiss	Q99466
别名	NOTCH4;INT3;notch 4

产品验证



实验步骤

访问官网浏览详情: www.ablybio.cn