

TRIB3 Rabbit mAb

货号: B31165

产品信息

| 反应 | Human |
|-------|--------------------------------------------------------------------------------------------|
| 宿主 | Rabbit |
| 克隆性 | Monoclonal |
| 预测反应 | |
| 应用 | WB IP |
| 推荐浓度 | WB: 1:500 - 1:2000 IP: 1:20 - 1:50 |
| 理论分子量 | 40kDa |
| 实测分子量 | 40kDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | K-562 |
| 细胞定位 | cytosol,nucleoplasm,nucleus,plasma membrane |
| 纯化 | Affinity purification |

抗原信息


| 抗原信息 | Recombinant fusion protein corresponding to Human TRIB3. |
|------|----------------------------------------------------------|
| 序列 | |

靶点信息

| 研究背景 | The protein encoded by this gene is a putative protein kinase that is induced by the transcription factor N F-kappaB. The encoded protein is a negative regulator of NF-kappaB and can also sensitize cells to TNF- a nd TRAIL-induced apoptosis. In addition, this protein can negatively regulate the cell survival serine-threo nine kinase AKT1. Differential promoter usage and alternate splicing result in multiple transcript variants. [provided by RefSeq, Jul 2014] |
|--------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| 基因 ID | 57761 |

| 基因名 | TRIB3 |
|-------|-----------------------------------|
| Swiss | Q96RU7 |
| 别名 | C20orf97; NIPK; SINK; SKIP3; TRB3 |

产品验证



Western blot analysis of TRIB3 expressed in K-562 using TRIB3 Rabbit mAb at 1:1000. Secondary antibod y: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn